Fluid Phase Equilibria, 68 (1991) 297–298
Elsevier Science Publishers B.V., Amsterdam

Contents Volume 68 (1991)

Volume 68

Correlation of vapour–liquid equilibria for tautomeric mixtures of acetylacetone with non-associated components
I. Nagata and Y. Fukushima (Kanazawa, Japan) 1–12
Calculation of heats of vaporization of halogenated hydrocarbons from saturated vapour pressure data
P. Basařová and V. Svoboda (Prague, Czechoslovakia) 13–34
Influence of size and shape effects on the solubility of hydrocarbons: the role of the combinatorial entropy
K. Kniaź (Warsaw, Poland) 35–46
Group-contribution equation of state for correlating and predicting thermodynamic properties of weakly polar and non-associating mixtures. Binary and multicomponent systems
W. Abdoul, E. Rauzy and A. Péneloux (Marseille, France) 47–102
Correlation of the solubility of hexacosane in aliphatic alcohols
U. Domańska and K. Domański (Warsaw, Poland) 103–114
Behavior of the LEMF equation for vapor–liquid equilibrium data treatment of systems with negative deviations from ideality
R.B. Mato, F.A. Mato and F. Mato (Valladolid, Spain) 115–130
Measurement of infinite dilution activity coefficients using high performance liquid chromatography
I.G. Economou, P. Vimalchand and M.D. Donohue (Baltimore, MD, USA) . 131–149
Thermodynamics of binary mixtures containing organic carbonates. Part VI. Isothermal vapor–liquid equilibria for dimethyl carbonate + normal alkanes
M.J. Cocero, I. García, J.A. González and J.C. Cobos (Valladolid, Spain) . . 151–161
A modified method for vapour–liquid equilibria measurement by inert gas stripping
J. Ovečková, J. Surový and E. Graczová (Bratislava, Czechoslovakia) 163–172
A new variant of the Rayleigh distillation method for the determination of limiting activity coefficients
V. Dohnal and I. Horáková (Prague, Czechoslovakia) 173–185
Liquid–liquid miscibility for binary systems: *N*-methylpyrrolidone + *n*-alkane and propanenitrile + *n*-alkane
R. Eustaquio-Rincón, R. Molnar and A. Trejo (México, México) 187–195
Solubilities of nonane vapor in styrene–butadiene copolymers at 100 and 130 °C
Y. Iwai, T. Ishidao, S. Miyamoto, H. Ikeda and Y. Arai (Fukuoka, Japan) . . 197–205
Prediction of solid–liquid equilibria from measured adduction constants
X. Feng, S.-K. Wang and J. Shi (Jiangsu, People's Republic of China) 207–218
Vapour–liquid equilibrium in the binary system ethanoic acid (1)–cyclohexane (2) at 101.325 kPa
I. Malijevská, K. Chýlková and M. Sýsová (Prague, Czechoslovakia) 219–228
Vapor–liquid equilibrium of the system H_2SO_4–SO_3. Part I. Vapor pressure measurements with a new static vapor pressure apparatus
J. Schrage (Aachen, Germany) 229–245

Vapor–liquid equilibrium of the system H_2SO_4–SO_3. Part II. Thermodynamic description with regard to the formation of $H_2S_2O_7$
J. Nilges and J. Schrage (Aachen, Germany) 247–261
Solubilities of theobromine and caffeine in supercritical carbon dioxide: correlation with density-based models
S. Li, G.S. Varadarajan and S. Hartland (Zurich, Switzerland) 263–280
Vapour pressures for binary liquid mixtures of methylethylketone with methylene chloride, 1,2-dichloroethane, trichloroethene, tetrachloroethene and cyclohexane
J.R. Nath and R. Saini (Gorakhpur, India) 281–290

Announcements 291

Erratum 293

Author Index 295

Contents of Volume 68 (1991) 297–298